For Research Use Only

TRPA1 Polyclonal antibody

Catalog Number:25226-1-AP




Basic Information	Catalog Number: 25226-1-AP	GenBank Accession Number: BC 153003	Purification Method: Antigen affinity purification	
	Size: 150ul , Concentration: 500 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG18685	GeneID (NCBI): 8989 UNIPROT ID: 075762 Full Name: transient receptor potential cation channel, subfamily A, member 1 Calculated MW: 1119 aa, 128 kDa Observed MW: 130 kDa	Recommended Dilutions: WB 1:1000-1:8000	
Applications	Tested Applications: WB, ELISA Species Specificity: human		Positive Controls: WB : A549 cells, U2OS cells	
Background Information	TRPA1 (Transient Receptor Potential Ankyrin 1) is a member of the transient receptor potential (TRP) channel family and is a non-selective cation channel. TRPA1 is predominantly expressed in sensory neurons and is involved in a variety of physiological activities, such as stabilizing cellular membrane potentials, maintaining cellular fluid homeostasis, and regulating intercellular signaling. TRPA1 is a multimodal cell membrane receptor that senses different stimuli and upon activation generates action potential signals via osmolality, temperature and inflammatory factors. TRPA1 has been associated with neurogenic inflammation, osteoarthritis, atopic dermatitis, asthma, inflammatory bowel disease, migraine headache, cancer pain, gout and other diseases, and it is a promising target for drug therapy.			
Storage *** 20ul sizes contain 0.1% BSA	Storage: Store at -20°C. Stable for one year aff Storage Buffer: PBS with 0.02% sodium azide and 50 Aliquoting is unnecessary for -20°C s	9% glycerol pH 7.3.		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 25226-1-AP (TRPA1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.