For Research Use Only

## FER Polyclonal antibody Catalog Number: 25287-1-AP 2 Publications




Basic Information	Catalog Number: 25287-1-AP	GenBank Accession Number: BC017060	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 800 ug/ml by	2241	WB 1:500-1:2000	
	Nanodrop and 433 ug/ml by Bradford	UNITROTTE.	IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	method using BSA as the standard; Source:	P16591	protein lysate IHC 1:50-1:500	
	Rabbit	Full Name: fer (fps/fes related) tyrosine kinase	IF/ICC 1:20-1:200	
	Isotype: IgG	Calculated MW: 822 aa, 95 kDa		
	Immunogen Catalog Number: AG18173	Observed MW: 95 kDa		
Applications	Tested Applications: Positive Controls:			
	WB, IHC, IF/ICC, IP, ELISA	WB:NIH/3	WB : NIH/3T3 cells, A431 cells, HeLa cells, MCF-7 cells	
	Cited Applications: IP : MCF-7 ce WB Species Specificity: IHC : human		cells,	
			n kidney tissue, human colon cancer tissue	
	human, mouse	IF/ICC : MC	F-7 cells,	
	Cited Species: human			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	retrieval may be performed w			
Background Information	retrieval may be performed w buffer pH 6.0 FER(Tyrosine-protein kinase Fer) is a protein kinase acts downstream of ce	<b>ith citrate</b> lso named as TYK3 and belongs to t ll surface receptors for growth facto nbly, lamellipodia formation, cell a	ne protein kinase superfamily. Tyrosine- rs and plays a role in the regulation of the Idhesion, cell migration and chemotaxis. alternative splicing.	
	retrieval may be performed w buffer pH 6.0 FER(Tyrosine-protein kinase Fer) is a protein kinase acts downstream of ce actin cytoskeleton, microtubule asser This protein has 3 isoforms produced	<b>ith citrate</b> lso named as TYK3 and belongs to t ll surface receptors for growth facto nbly, lamellipodia formation, cell a	rs and plays a role in the regulation of the adhesion, cell migration and chemotaxis.	
Background Information Notable Publications	retrieval may be performed w buffer pH 6.0 FER(Tyrosine-protein kinase Fer) is a protein kinase acts downstream of ce actin cytoskeleton, microtubule asser This protein has 3 isoforms produced Author Pub	<b>ith citrate</b> Iso named as TYK3 and belongs to t Il surface receptors for growth facto mbly, lamellipodia formation, cell a by alternative promoter usage and	rs and plays a role in the regulation of the adhesion, cell migration and chemotaxis. alternative splicing.	
	retrieval may be performed w buffer pH 6.0FER(Tyrosine-protein kinase Fer) is a protein kinase acts downstream of ce actin cytoskeleton, microtubule asser This protein has 3 isoforms producedAuthorPub Xue YangXue Yang322	ith citrate Iso named as TYK3 and belongs to t Il surface receptors for growth facto mbly, lamellipodia formation, cell a by alternative promoter usage and med ID Journal	rs and plays a role in the regulation of the adhesion, cell migration and chemotaxis. alternative splicing. Application	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 25287-1-AP (FER Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




IP result of anti-FER (IP:25287-1-AP, 4ug; Detection:25287-1-AP 1:1000) with MCF-7 cells lysate 2400ug.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 25287-1-AP (FER antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 25287-1-AP (FER antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of MCF-7 cells using 25287-1-AP (FER antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).