For Research Use Only

CEBPZ Polyclonal antibody

Catalog Number: 25612-1-AP 2 Publications



Basic Information

Catalog Number:

GenBank Accession Number: BC034475

25612-1-AP

GeneID (NCBI): Size: 150ul , Concentration: 450 ug/ml by 10153

Nanodrop; **UNIPROT ID:** Q03701

Rabbit Full Name:

Isotype: CCAAT/enhancer binding protein (C/EBP), zeta

IgG Immunogen Catalog Number: Calculated MW: 1054 aa, 121 kDa AG11003

> Observed MW: 120-130 kDa

Applications

Tested Applications: WB, IF/ICC, ELISA

Cited Applications:

Species Specificity: human, mouse **Cited Species:** human, mouse

Positive Controls:

WB: HeLa cells, mouse testis tissue, Jurkat cells, K-562

Purification Method:

WB 1:1000-1:6000 IF/ICC 1:20-1:200

Antigen affinity purification

Recommended Dilutions:

cells

IF/ICC: HeLa cells,

Background Information

CEBPZ (CCAAT/enhancer binding protein zeta) belongs to the CBF/Mak21 family, it plays a role in cellular response to environmental stimuli through a transcriptional process that involves heat shock factors, conserved DNA elements (heat shock elements or HSEs) and CCAAT boxes. The protein acts as a DNA-binding transcriptional activator and regulates the heat-shock protein 70 (HSP70) promoter in a CCAAT-dependent manner.

Notable Publications

Author	Pubmed ID	Journal	Application
Hanxiao Shi	33934437	Cancer Sci	WB
Tong Wu	34478875	Mol Cell Proteomics	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 25612-1-AP (CEBPZ Antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using 25612-1-AP (CEBPZ antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 25612-1-AP (CEBPZ antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 25612-1-AP (CEBPZ antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.