

For Research Use Only

UHRF2 Polyclonal antibody

Catalog Number: 25710-1-AP **1 Publications**



Basic Information

Catalog Number: 25710-1-AP	GenBank Accession Number: BC028397	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 300 µg/ml by Nanodrop;	GeneID (NCBI): 115426	Recommended Dilutions: WB 1:500-1:2000 IP 0.5-4.0 ug for IP and 1:200-1:1000 for WB
Source: Rabbit	Full Name: ubiquitin-like with PHD and ring finger domains 2	
Isotype: IgG	Calculated MW: 90 kDa	
Immunogen Catalog Number: AG22519	Observed MW: 90 kDa	

Applications

Tested Applications: IP, WB, ELISA	Positive Controls: WB : HL-60 cells, HeLa cells, Jurkat cells, PC-3 cells IP : Jurkat cells,
Cited Applications: WB	
Species Specificity: human	
Cited Species: human	

Background Information

Ubiquitin-Like with PHD and ring finger domains 2 (UHRF2), a member that belongs to the family of UHRF, contains five recognizable functional domains, namely the ubiquitin-like domain (UBL) domain, tandem-Tudor domain (TTD), plant homeodomain (PHD), SET and RING associated (SRA) domain, and really interesting new gene (RING) finger domain. Due to the complex structure, UHRF2 possesses multiple functions in diverse cellular processes. As a ubiquitin E3 ligase, UHRF2 could ubiquitinate PCNP, a nuclear protein that contains two remarkable PEST sequences which are rich in proline (P), glutamic acid (E), serine (S), and threonine (T). It has been also reported that UHRF2 could serve as a vital cell cycle regulator by interacting with multiple cyclins, CDKs, p53, pRB and PCNA. UHRF2 has been revealed to possess epigenetic regulation function and is capable of maintaining 5mC levels in certain genomic loci in brain and stabilizes TIP60 to regulate H3K9ac and H3K14ac through RING finger domain. Moreover, UHRF2 could promote DNA damage repair by reducing the level of p21 mediated by RING finger domain. Recently, emerging evidence indicated that UHRF2 was involved in the tumorigenesis and progression of several human cancers, such as esophageal squamous cell carcinoma, lung cancer and colorectal cancer. UHRF2 has 2 isoforms with the molecular mass of 56 and 90 kDa. (PMID: 34400880)

Notable Publications

Author	Pubmed ID	Journal	Application
Shengjun Geng	35732617	Cell Death Dis	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 25710-1-AP (UHRF2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP Result of anti-UHRF2 (IP:25710-1-AP, 4ug; Detection:25710-1-AP 1:300) with Jurkat cells lysate 4000ug.