

For Research Use Only

ROR2 Polyclonal antibody

Catalog Number: 25842-1-AP



Basic Information

Catalog Number:

25842-1-AP

Size:

150ul , Concentration: 700 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG22959

GenBank Accession Number:

BC130522

GeneID (NCBI):

4920

UNIPROT ID:

Q01974

Full Name:

receptor tyrosine kinase-like orphan receptor 2

Calculated MW:

943 aa, 105 kDa

Observed MW:

105 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:8000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human

Positive Controls:

WB : HeLa cells, HepG2 cells, T-47D cells, K-562 cells

Background Information

Tyrosine-protein kinase transmembrane receptor ROR2 belongs to the ROR-family of receptor tyrosine kinases (RTKs) (PMID: 12815588). ROR2 is a developmentally regulated protein that has significant expression and roles in a wide variety of tissues during early embryonic development. It was established that ROR2 plays a key role in skeletal and neural organogenesis, as well as in midgut development in the mouse, but then becomes repressed in adult tissues. It may act as a receptor for wnt ligand WNT5A which may result in the inhibition of WNT3A-mediated signaling (PMID: 25614331). Mutations in ROR2 gene are associated with autosomal recessive Robinow syndrome and autosomal dominant brachydactyly type B (PMID: 12815588).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 25842-1-AP (ROR2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.