

For Research Use Only

C9 Polyclonal antibody

Catalog Number: 25891-1-AP



Basic Information

Catalog Number: 25891-1-AP	GenBank Accession Number: BC020721	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 500 ug/ml by Nanodrop and 493 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 735	Recommended Dilutions: WB 1:1000-1:8000
Source: Rabbit	UNIPROT ID: P02748	
Isotype: IgG	Full Name: complement component 9	
Immunogen Catalog Number: AG23137	Calculated MW: 559 aa, 63 kDa	
	Observed MW: 64 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : human plasma,
Species Specificity: human	

Background Information

The final consequence of complement activation is the formation of the membrane attack complex (MAC), a multiprotein assembly that perforates cell membranes, forming transmembrane channels. The MAC plays a critical role in complement-mediated hemolysis and inflammasome activation (PMID: 27444648). Complement component 9 (C9) is 1 of 5 late-acting proteins of complement that participate in the assembly and function of the MAC. C9 may be cleaved by thrombin to generate fragments C9a (25 kDa) and C9b (37 kDa) (PMID: 7523406).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human plasma were subjected to SDS PAGE followed by western blot with 25891-1-AP (C9 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.