## For Research Use Only

## C9 Polyclonal antibody

Catalog Number: 25891-1-AP



**Purification Method:** 

WB 1:1000-1:8000

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number: GenBank Accession Number: 25891-1-AP BC020721

Size: GeneID (NCBI):

150ul , Concentration: 500 µg/ml by 735

Nanodrop and 493 µg/ml by Bradford Full Name:

method using BSA as the standard; complement component 9

Source: Calculated MW:
Rabbit 559 aa, 63 kDa
Isotype: Observed MW:
IgG 64 kDa

Immunogen Catalog Number:

AG23137

Tested Applications: Positive Controls:

WB, ELISA WB: human plasma,

Species Specificity:

human

## **Background Information**

The final consequence of complement activation is the formation of the membrane attack complex (MAC), a multiprotein assembly that perforates cell membranes, forming transmembrane channels. The MAC plays a critical role in complement-mediated hemolysis and inflammasome activation (PMID: 27444648). Complement component 9 (C9) is 1 of 5 late-acting proteins of complement that participate in the assembly and function of the MAC. C9 may be cleaved by thrombin to generate fragments C9a (25 kDa) and C9b (37 kDa) (PMID: 7523406).

Storage

**Applications** 

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## **Selected Validation Data**



human plasma were subjected to SDS PAGE followed by western blot with 25891-1-AP (C9 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.