For Research Use Only

PUM1 Polyclonal antibody

Catalog Number:26256-1-AP

Featured Product

2 Publications



Basic Information

Catalog Number:

26256-1-AP

Size: 150ul, Concentration: 500 µg/ml by

Nanodrop and 307 µg/ml by Bradford Full Name:

method using BSA as the standard;

Rabbit Isotype: IgG

Immunogen Catalog Number:

AG24651

GenBank Accession Number:

BC013398

GeneID (NCBI): 9698

pumilio homolog 1 (Drosophila)

Calculated MW: 1186 aa, 126 kDa

Observed MW:

130-140 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:4000

IP 0.5-4.0 ug for IP and 1:200-1:1000

for WB IF 1:50-1:500

Applications

Tested Applications: IF, IP, WB, ELISA

Cited Applications:

IHC, WB

Species Specificity:

human

Cited Species:

human

Positive Controls:

WB: MCF-7 cells, IP: MCF-7 cells,

IF: MCF-7 cells,

Background Information

PUM1, also named as Pumilio homolog 1, is a 1186 amino acid protein, which is expressed in brain, heart, kidney, muscle, intestine and stomach. PUM1 as sequence-specific RNA-binding protein that acts as a post-transcriptional repressor by binding the 3'-UTR of mRNA targets. PUM1 binds to an RNA consensus sequence, the Pumilio Response Element (PRE), 5'-UGUANAUA-3', that is related to the Nanos Response Element (NRE). The calculated molecular weight of PUM1 is 126 kDa, but the modified PUM1 is about 130-140 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Beatriz Escudero-Paniagua	31095674	Carcinogenesis	WB
Xue Guan	29428722	Biochem Biophys Res Commun	WB,IHC

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



MCF-7 cells were subjected to SDS PAGE followed by western blot with 26256-1-AP (PUM1 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



IP Result of anti-PUM1 (IP:26256-1-AP, 4ug; Detection:26256-1-AP 1:300) with MCF-7 cells lysate 3200ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using 26256-1-AP (PUM1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).