For Research Use Only

## DALRD3 Polyclonal antibody

Catalog Number: 26294-1-AP 1 Publications





Basic Information	Catalog Number: 26294-1-AP			Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500	
	Size: 150ul , Concentration: 1000 ug/ml by Nanodrop and 393 ug/ml by Bradford method using BSA as the standard;				
	Source: Rabbit	Full Name: DALR anticodon b	ainding domain		
	lsotype: IgG	containing 3 Observed MW:			
	Immunogen Catalog Number: AG23437	er: 58 kDa			
Applications	Tested Applications: WB, IHC, EUSA		Positive Cont		
	Cited Applications: WB	WB : K-562 cells, COLO 320 cells IHC : human colon cancer tissue,			
	Species Specificity: human				
	Cited Species: human				
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen			
Notable Publications	Author Pub	omed ID Jo	ournal		Application
	Jenna M Lentini 324	27860 N	at Commun		WB
Storage	Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50				

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data






COLO 320 cells were subjected to SDS PAGE followed by western blot with 26294-1-AP (DALRD3 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 26294-1-AP (DALRD3 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 26294-1-AP (DALRD3 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



K-562 cells were subjected to SDS PAGE followed by western blot with 26294-1-AP (DALRD3 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.