For Research Use Only

SMARCA2-Specific Polyclonal antibody



Catalog Number: 26613-1-AP

Featured Product

4 Publications

Basic Information

Catalog Number: 26613-1-AP

150ul, Concentration: 700 µg/ml by

Nanodrop and 393 µg/ml by Bradford Full Name: method using BSA as the standard;

Rabbit

IgG Immunogen Catalog Number:

AG25157

Isotype:

Size:

GenBank Accession Number:

NM 003070 GeneID (NCBI):

SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 2

Calculated MW: 181 kDa Observed MW: 180-190 kDa

Applications

Tested Applications:

IF, IHC, WB, ELISA **Cited Applications:**

Species Specificity: human, mouse **Cited Species:** human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000 IHC 1:50-1:500 IF 1:50-1:500

Positive Controls:

WB: HeLa cells, NIH/3T3 cells, MDA-MB-453s cells

IHC: human ovary tumor tissue,

IF: HepG2 cells,

Background Information

SMARCA2 is a transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. SMARCA2 is involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association via its associatcomplex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligandbound VDR-mediated transrepression of the CYP27B1 gene.

Notable Publications

Author	Pubmed ID	Journal	Application
Debolina Ganguly	30171737	Stem Cells	WB
Zhixiong Liu	30876848	Neuron	WB
Xiaoli Xu	30127991	Oncol Lett	WB

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 26613-1-AP (SMARCA2-Specific antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 26613-1-AP (SMARCA2-Specific antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 26613-1-AP (SMARCA2-Specific antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



WB result of SMARCA2-Specific antibody (26613-1-AP; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SMARCA2-Specific transfected HeLa cells.