For Research Use Only

## MYO6 Polyclonal antibody

Catalog Number:26778-1-AP

Featured Product


7 Publications




Basic Information	Catalog Number: 26778-1-AP	GenBank Accession BC146764	Number:	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):		Recommended Dilutions:	
	150ul , Concentration: 350 ug/ml by	4646		WB 1:5000-1:50000	
	Nanodrop and 333 ug/ml by Bradford	UNIPROT ID:		IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	method using BSA as the standard;	Q9UM54		protein lysate	
	Source:	Full Name:		IHC 1:50-1:500	
	Rabbit	myosin VI		IF-P 1:50-1:500 IF/ICC 1:200-1:800	
	Isotype:	Calculated MW:		17700 1.200 1.000	
	IgG	1285 aa, 149 kDa			
	Immunogen Catalog Number: AG24906	Observed MW: 145-150 kDa			
Applications	Tested Applications:		Positive Cont	rols:	
				WB : HEK-293 cells, mouse small intestine tissue, DI 145 cells, LNCaP cells, MCF-7 cells	
	WB, IHC, IF			•	
	Species Specificity:		IP: PC-3 cells		
	human, mouse		IHC : human p intestine tiss	rostate cancer tissue, human small Ie	
	Cited Species: human, mouse		IF-P: mouse s	mall intestine tissue,	
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen			
Background Information	MYO6, an actin-based motor protein, MYO6 is highly expressed in the inne kidney proximal tubule cells and inte within cells, including clathrin-media autophagy (PMID: 23620821; PMID: 23 of cancer, and it has been widely repo indicate that MYO6 is associated with (PMID: 29022908).	r and outer hair cells estinal enterocytes. A ated endocytosis, ves 8591580). Previous st orted to contribute to	of the ear, retina, nd it participates icular membrane udies showed tha tumor cell migra	and polarized epithelial cells such as in a wide range of biological process traffic, polarized secretion, and tt MYO6 is upregulated in various type ion and metastasis. Some articles	
	MYO6 is highly expressed in the inne kidney proximal tubule cells and inte within cells, including clathrin-media autophagy (PMID: 23620821; PMID: 22 of cancer, and it has been widely repo indicate that MYO6 is associated with (PMID: 29022908).	r and outer hair cells estinal enterocytes. A ated endocytosis, ves 8591580). Previous st orted to contribute to n prostate cancer, lung	of the ear, retina, nd it participates icular membrane udies showed tha tumor cell migra	and polarized epithelial cells such as in a wide range of biological process traffic, polarized secretion, and It MYO6 is upregulated in various type ion and metastasis. Some articles solorectal cancer and gastric cancer	
	MYO6 is highly expressed in the inner kidney proximal tubule cells and inter within cells, including clathrin-media autophagy (PMID: 23620821; PMID: 2 of cancer, and it has been widely repor indicate that MYO6 is associated with (PMID: 29022908). Author Pul	r and outer hair cells estinal enterocytes. A ated endocytosis, ves 8591580). Previous st orted to contribute to a prostate cancer, lung bmed ID Jou	of the ear, retina, nd it participates icular membrane udies showed tha tumor cell migra g cancer, human o	and polarized epithelial cells such as in a wide range of biological process traffic, polarized secretion, and tt MYO6 is upregulated in various type ion and metastasis. Some articles	
	MYO6 is highly expressed in the inner kidney proximal tubule cells and inter within cells, including clathrin-media autophagy (PMID: 23620821; PMID: 23 of cancer, and it has been widely report indicate that MYO6 is associated with (PMID: 29022908).AuthorPulShichen Hu31	r and outer hair cells estinal enterocytes. A sted endocytosis, ves 8591580). Previous st orted to contribute to a prostate cancer, lung bmed ID Jou 371777 Nat	of the ear, retina, nd it participates icular membrane udies showed tha tumor cell migra g cancer, human o	and polarized epithelial cells such as in a wide range of biological process traffic, polarized secretion, and it MYO6 is upregulated in various type ion and metastasis. Some articles colorectal cancer and gastric cancer Application WB	
	MYO6 is highly expressed in the inner kidney proximal tubule cells and inter within cells, including clathrin-media autophagy (PMID: 23620821; PMID: 20 of cancer, and it has been widely repr indicate that MYO6 is associated with (PMID: 29022908).AuthorPulShichen Hu31: Shengfeng Zheng39.	r and outer hair cells estinal enterocytes. A ated endocytosis, ves 8591580). Previous st orted to contribute to prostate cancer, lung bmed ID Jou 371777 Nat 449086 Cel	of the ear, retina, nd it participates icular membrane udies showed tha tumor cell migra g cancer, human o rnal Commun	and polarized epithelial cells such as in a wide range of biological process traffic, polarized secretion, and it MYO6 is upregulated in various type ion and metastasis. Some articles olorectal cancer and gastric cancer Application WB	
Background Information	MYO6 is highly expressed in the inner kidney proximal tubule cells and inter within cells, including clathrin-media autophagy (PMID: 23620821; PMID: 20 of cancer, and it has been widely repr indicate that MYO6 is associated with (PMID: 29022908).AuthorPulShichen Hu31: Shengfeng Zheng39.	r and outer hair cells estinal enterocytes. A sted endocytosis, ves 8591580). Previous st orted to contribute to a prostate cancer, lung bmed ID Jou 371777 Nat 449086 Cel 445095 Car er shipment. % glycerol pH 7.3.	of the ear, retina, nd it participates icular membrane udies showed tha tumor cell migra g cancer, human o rnal Commun I Commun Signa	and polarized epithelial cells such as in a wide range of biological process traffic, polarized secretion, and tt MYO6 is upregulated in various type ion and metastasis. Some articles solorectal cancer and gastric cancer Application WB I. IHC	

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 26778-1-AP (MYO6 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed mouse small intestine tissue using 26778-1-AP (MYO6 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using MYO6 antibody (26778-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).



IP result of anti-MYO6 (IP:26778-1-AP, 4ug: Detection:26778-1-AP 1:500) with PC-3 cells lysate 2000 ug.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 26778-1-AP (MYO6 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).