

For Research Use Only

MYO6 Polyclonal antibody

Catalog Number: 26778-1-AP **4 Publications**



Basic Information

Catalog Number: 26778-1-AP	GenBank Accession Number: BC146764	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 350 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 4646	Recommended Dilutions: WB 1:500-1:1000 IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
Source: Rabbit	Full Name: myosin VI	IHC 1:50-1:500 IF 1:50-1:500
Isotype: IgG	Calculated MW: 1285 aa, 149 kDa	
Immunogen Catalog Number: AG24906	Observed MW: 145-150 kDa	

Applications

Tested Applications: IF, IHC, IP, WB, ELISA	Positive Controls: WB : mouse small intestine tissue, IP : PC-3 cells, IHC : human prostate cancer tissue, human small intestine tissue IF : mouse small intestine tissue,
Cited Applications: IF, IHC, WB	
Species Specificity: human, mouse	
Cited Species: human, mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

MYO6, an actin-based motor protein, is the only myosin known to move toward the minus end of actin filaments. MYO6 is highly expressed in the inner and outer hair cells of the ear, retina, and polarized epithelial cells such as kidney proximal tubule cells and intestinal enterocytes. And it participates in a wide range of biological processes within cells, including clathrin-mediated endocytosis, vesicular membrane traffic, polarized secretion, and autophagy (PMID: 23620821; PMID: 28591580). Previous studies showed that MYO6 is upregulated in various types of cancer, and it has been widely reported to contribute to tumor cell migration and metastasis. Some articles indicate that MYO6 is associated with prostate cancer, lung cancer, human colorectal cancer and gastric cancer (PMID: 29022908).

Notable Publications

Author	Pubmed ID	Journal	Application
Shichen Hu	31371777	Nat Commun	WB
Xubo Chen	37859596	Int J Toxicol	IF
He Cao	37794006	Nat Commun	WB, IHC, IF

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




mouse small intestine tissue were subjected to SDS PAGE followed by western blot with 26778-1-AP (MYO6 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed mouse small intestine tissue using 26778-1-AP (MYO6 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-MYO6(IP:26778-1-AP, 4ug; Detection:26778-1-AP 1:500) with PC-3 cells lysate 2000 ug.



Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 26778-1-AP (MYO6 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).