## For Research Use Only

## YTHDF1 Polyclonal antibody

Catalog Number: 26787-1-AP 5 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number: BC050284

26787-1-AP GeneID (NCBI):

150ul , Concentration: 300 ug/ml by 54915

Bradford method using BSA as the

standard;

Size:

Source:

Rabbit YTH domain family, member 1

Isotype: Calculated MW: 559 aa, 61 kDa

Immunogen Catalog Number: Observed MW:

AG25347

Positive Controls:

WB: HepG2 cells,

IP: HepG2 cells,

60 kDa

**UNIPROT ID:** 

Full Name:

Q9BYJ9

**Applications** 

**Tested Applications:** 

WB, IP, ELISA

**Cited Applications:** 

WB, IHC, RIP

Species Specificity: human

**Cited Species:** 

human, mouse

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000

**Purification Method:** 

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

**Background Information** 

YTHDF1, also named YTH domain-containing family protein 1 or C20orf21, is a 559 amino acid protein, which localizes in the cytoplasm. YTHDF1 specifically recognizes and binds N6-methyladenosine (m6A)-containing mRNAs, and promotes mRNA translation efficiency. M6A is a modification present at the internal sites of mRNAs and  $some\ non-coding\ RNAs\ and\ plays\ a\ role\ in\ the\ efficiency\ of\ mRNA\ splicing,\ processing,\ and\ stability.\ YTHDF1\ acts\ as$  $a \ regulator \ of \ mRNA \ translation \ efficiency: promotes \ ribosome \ loading \ to \ m6A-containing \ mRNAs \ and \ interacts \ with$ translation initiation factors eIF3 (EIF3A or EIF3B) to facilitate translation initiation. YTHDF1 exists two isoforms and the calculated molecular weight of isoforms are 61 kDa and 21 kDa.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Xiaohui Bai	35939138	Cell Mol Neurobiol	WB,IHC
Chaopeng Ou	39304892	BMC Med	WB,IHC
Danyu Chen	38900944	Cancer Res	WB,RIP,IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 26787-1-AP (YTHDF1 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP result of anti-YTHDF1 (IP:26787-1-AP, 4ug; Detection:26787-1-AP 1:300) with HepG2 cells lysate 3200ug.