

For Research Use Only

BCLAF1 Polyclonal antibody

Catalog Number: 26809-1-AP

Featured Product

8 Publications



Basic Information

Catalog Number: 26809-1-AP	GenBank Accession Number: BC132780	Purification Method: Antigen affinity purification
Size: 150ul, Concentration: 400 ug/ml by Nanodrop and 267 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 9774	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:250-1:1000
Source: Rabbit	UNIPROT ID: Q9NYF8	
Isotype: IgG	Full Name: BCL2-associated transcription factor 1	
Immunogen Catalog Number: AG25210	Calculated MW: 920 aa, 106 kDa	
	Observed MW: 145 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls: WB : Jurkat cells, MCF-7 cells IHC : mouse testis tissue, human colon tissue
Cited Applications: WB, IHC, IF, RIP	
Species Specificity: human, mouse	
Cited Species: human, mouse, pig	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

BCLF1, also named as Bcl-2-associated transcription factor 1, is a 920 amino acid protein, which Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA. BCLF1 as a death-promoting transcriptional repressor may be involved in cyclin-D1/CCND1 mRNA stability through the SNARP complex which associates with both the 3'end of the CCND1 gene and its mRNA. The calculated molecular weight of BCLF1 is a 110 kDa, but the modified BCLF1 protein is about 140-150 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Wenchong Tan	36137350	Transl Oncol	WB,IHC,IF
Mingwei Chen	33188158	Aging (Albany NY)	WB
Wenbo Liu	36355419	eLife	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 26809-1-AP (BCLAF1 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 26809-1-AP (BCLAF1 antibody) at dilution of 1:500 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 26809-1-AP (BCLAF1 antibody) at dilution of 1:500 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 26809-1-AP (BCLAF1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 26809-1-AP (BCLAF1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).