

PLEKHA1 Polyclonal antibody

Catalog Number: 26830-1-AP

Basic Information

Catalog Number: 26830-1-AP	GenBank Accession Number: BC001136	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 750 ug/ml by Nanodrop and 400 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 59338	Recommended Dilutions: WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500 IF/ICC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q9HB21	
Isotype: IgG	Full Name: pleckstrin homology domain containing, family A (phosphoinositide binding specific) member 1	
Immunogen Catalog Number: AG25285	Calculated MW: 46 kDa	
	Observed MW: 46 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA	Positive Controls:
Species Specificity: human	WB : HeLa cells, A549 cells, Jurkat cells, MOLT-4 cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IP : A549 cells,
	IHC : human colon tissue, human pancreas tissue
	IF/ICC : A549 cells,

Background Information

Pleckstrin homology (PH) domain is commonly found in eukaryotic signaling proteins and possesses multiple functions including the abilities to bind inositol phosphates and various proteins. The tandem PH domain containing protein-1 (TAPP1) or PH domain containing-family A (phosphoinositide binding specific) member 1 (PLEKHA1), interacts strongly and specifically with phosphatidylinositol 3,4-trisphosphate [PtdIns(3,4)P(2)], which is one of the immediate breakdown products of PtdIns(3,4,5) P (3) and functions as a signalling molecule in insulin- and growth-factor-stimulated pathways. TAPP1 is also associated with the protein- tyrosine-phosphatase-like protein-1 (PTPL1 also known as FAP-1) and maintains PTPL1 in cytoplasm. By binding to PtdIns(3,4) P (2) and PTPL1, TAPP1 may regulate the membrane localization of PTPL1."

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 26830-1-AP (PLEKHA1 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 26830-1-AP (PLEKHA1 antibody) at dilution of 1:200 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 26830-1-AP (PLEKHA1 antibody) at dilution of 1:200 (under 40x lens).



Various lysates were subjected to SDS PAGE followed by western blot with 26830-1-AP (PLEKHA1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-PLEKHA1 (IP:26830-1-AP, 4ug; Detection:26830-1-AP 1:1000) with A 549 cells lysate 1600 ug.



Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using PLEKHA1 antibody (26830-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).