For Research Use Only

JAM-A/CD321 Polyclonal antibody

Catalog Number:26941-1-AP 1 Publications



Basic Information

Catalog Number: 26941-1-AP

Size:

Isotype

GenBank Accession Number:

BC001533

GeneID (NCBI):

150ul , Concentration: 600 ug/ml by Nanodrop and 433 ug/ml by Bradford $\,$ UNIPROT ID: method using BSA as the standard;

50848 Q9Y624

Source: Rabbit

Full Name: F11 receptor Calculated MW:

Immunogen Catalog Number:

Observed MW: AG25641

33 kDa

33 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:4000 IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB

Species Specificity:

human **Cited Species:**

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HUVEC cells, human peripheral blood platelets,

U-87 MG cells

IHC: human breast cancer tissue, human spleen tissue

Background Information

The F11 receptor (F11R) was first identified in human platelets as a 32/35 kDa protein duplex that serves as the receptor for a functional antibody that activates platelets. It seems to play a role in epithelial tight junction formation. It was also reported to function as a receptor for reovirus and ligand for the integrin LFA1.

Notable Publications

Author **Pubmed ID** Journal Application WB Qianyao Zhang 39956317 Neuropharmacology

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 26941-1-AP (JAM-A/Junctional Adhesion Molecule A antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 26941-1-AP (JAM-A/Junctional Adhesion Molecule A antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 26941-1-AP (JAM-A/Junctional Adhesion Molecule A antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.