

For Research Use Only

VPS37C Polyclonal antibody

Catalog Number: 27128-1-AP



Basic Information

Catalog Number: 27128-1-AP	GenBank Accession Number: BC141827	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 450 ug/ml by Nanodrop;	GeneID (NCBI): 55048	Recommended Dilutions: WB 1:500-1:2000
Source: Rabbit	UNIPROT ID: A5D8V6	IHC 1:50-1:500
Isotype: IgG	Full Name: vacuolar protein sorting 37 homolog C (S. cerevisiae)	IF/ICC 1:200-1:800
Immunogen Catalog Number: AG26024	Observed MW: 39-45 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls:
Species Specificity: human	WB : HT-29 cells, SK-BR-3 cells, T-47D cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : mouse colon tissue, mouse lung tissue
	IF/ICC : Raji cells,

Background Information

VPS37C (Vacuolar protein sorting-associated protein 37C) is also named as hVps37C, PML39 and ESCRT-I complex subunit VPS37C. VPS37C is a core subunit of the Endosomal sorting complex required for transport-I (ESCRT-I) complex (PMID: 37307706). VPS37C interacts with EKLF and prevents K48-linked polyubiquitination of EKLF and proteasome-mediated EKLF degradation, thus enhancing EKLF protein stability and transcriptional activity (PMID: 37307706).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 27128-1-AP (VP537C antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse colon tissue slide using 27128-1-AP (VP537C antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using 27128-1-AP (VP537C antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Raji cells using VP537C antibody (27128-1-AP) at dilution of 1:400 and Multi-rAb Coralite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).