

For Research Use Only

ASIC1 Polyclonal antibody

Catalog Number: 27235-1-AP

Featured Product

13 Publications



Basic Information

Catalog Number:

27235-1-AP

Size:

150ul, Concentration: 900 ug/ml by Nanodrop and 500 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG25941

GenBank Accession Number:

NM_001095

GeneID (NCBI):

41

UNIPROT ID:

P78348

Full Name:

amiloride-sensitive cation channel 2, neuronal

Calculated MW:

60 kDa

Observed MW:

60-70 kDa, 80-90 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

Applications

Tested Applications:

WB, IP, IHC, ELISA

Cited Applications:

WB, IHC, IF, CoIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse brain tissue, mouse cerebellum tissue, rat brain tissue

IP : mouse brain tissue,

IHC : human pancreas cancer tissue, mouse brain tissue

Background Information

ASIC1 (Acid-sensing ion channel 1), also named as ACCN2, BNaC2 or amiloride-sensitive cation channel 2, is a member of the ASICs family. ASICs, an H⁺-gated subgroup of the epithelial Na⁺ channel/ degenerin (ENaC/DEG) superfamily, are widely expressed throughout the central and peripheral nervous system and triggered by increased extracellular acidification (PMID: 28518134, 27941930). ASIC1 is a key player in acidosis-induced tumor growth and invasiveness. Clinically, alterations of ASIC1 are associated with poor survival in breast cancer (PMID: 26686084).

Notable Publications

Author	Pubmed ID	Journal	Application
Yangyang Li	36162458	Eur J Pharmacol	WB, IHC
Na Du	36155179	Life Sci	WB, IHC, IF
Wei Wang	35894166	Endocrinology	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 27235-1-AP (ASIC1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 27235-1-AP (ASIC1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-ASIC1 (IP:27235-1-AP, 4ug; Detection:27235-1-AP 1:300) with mouse brain tissue lysate 3000ug.



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 27235-1-AP (ASIC1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).