

For Research Use Only

# KMT2D Polyclonal antibody

Catalog Number: 27266-1-AP

Featured Product

9 Publications



## Basic Information

<b>Catalog Number:</b> 27266-1-AP	<b>GenBank Accession Number:</b> NM_003482	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 900 µg/ml by Nanodrop and 433 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 8085	<b>Recommended Dilutions:</b> WB 1:2000-1:16000 IHC 1:50-1:500
<b>Source:</b> Rabbit	<b>Full Name:</b> myeloid/lymphoid or mixed-lineage leukemia 2	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 593 kDa	
<b>Immunogen Catalog Number:</b> AG26195	<b>Observed MW:</b> 593 kDa	

## Applications

### Tested Applications:

IHC, WB, ELISA

### Cited Applications:

ChIP, IF, IHC, WB

### Species Specificity:

human

### Cited Species:

human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

WB : HEK-293 cells,

IHC : human breast cancer tissue, human colon cancer tissue, human lymphoma tissue

## Background Information

KMT2D (a COMPASS/ Set1 family member; also known as MLL4, ALR, and MLL2) is the biggest H3K4 methyltransferase among the most frequently mutated genes in many different types of cancer (PMID: 34194626). The enzymatic function of KMT2D depends on a cluster of C-terminal conserved domains, including a PHD domain, two FY-rich motifs (FYRC and FYRN) and a catalytic SET domain. KMT2D has two frequently occurring genomic alterations: truncation and missense mutations. Loss of KMT2D is a bona fide tumor suppressor gene in FL and DLBCL. Kmt2d perturbed the expression of genes that sustain proliferation and survival, and the KMT2D protein directly binds and associates with an active chromatin conformation in negative modulators of the BCR and lymphocyte migration pathways, which in turn could affect B cell responses to antigen (PMID: 26366712).

## Notable Publications

Author	Pubmed ID	Journal	Application
Xuejiao Leng	32879445	Oncogene	IHC
Ning Jiang	33200540	Thorac Cancer	IHC
Qianshuo Liu	34127651	Cell Death Discov	ChIP

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 27266-1-AP (KMT2D antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 27266-1-AP (KMT2D Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 27266-1-AP (KMT2D Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).