## For Research Use Only

## TRIM55 Polyclonal antibody

Catalog Number: 27274-1-AP



**Basic Information** 

Catalog Number: GenBank Accession Number:

27274-1-AP BC007750 GeneID (NCBI): Size: 150ul , Concentration: 700 µg/ml by 84675

Nanodrop and 433  $\mu$ g/ml by Bradford UNIPROT ID: method using BSA as the standard; Q9BYV6

Source: Full Name:

Rabbit tripartite motif-containing 55

Isotype: Calculated MW: 548 aa, 60 kDa Immunogen Catalog Number: Observed MW: AG25338 60-65 kDa 75-80 kDa **Purification Method:** Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000 IHC 1:50-1:500 IF 1:200-1:800

**Applications** 

Positive Controls: **Tested Applications:** IF, IHC, WB, ELISA WB: C2C12 cells,

Species Specificity: IHC: human liver cancer tissue, human heart tissue human, mouse, rat

IF: HeLa cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

## **Background Information**

Tripartite motif-containing 55 (TRIM55), which is also called muscle-specific ring finger 2 (Murf2), has been indicated to participate in the natural development of muscle and the cardiovascular system. The expression of TRIM55 also inhibited several markers of cancer stem-like cells (PMID: 34974792). MURF2 mRNA studies revealed up to four potential isoforms generated by differential splicing events within the 3'-region of the gene: p60, p50, p27, and p60B MURF2 isoforms. The change in the ratio of E3 ligase isoforms regulates the sequential activation of proteasomal or autophagic degradation of target components (PMID: 15199100, PMID: 24124537). This antibody could detect multiple isoforms or post-translational modifications of TRIM55.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


\*\*\* 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 27274-1-AP (TRIM55 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 27274-1-AP (TRIM55 antibody) at dilution of 1:100 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using TRIM55 antibody (27274-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).