For Research Use Only

JAK1 Polyclonal antibody Catalog Number:27284-1-AP 1 Publications



| Basic Information | Catalog Number: 27284-1-AP | GenBank Accession Number: BC132729 | Purification Method: Antigen affinity purification | |
|---------------------------------|--|---------------------------------------|--|--|
| | Size: 150ul, Concentration: 850 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG25579 | GenelD (NCBI): | Recommended Dilutions: WB: 1:500-1:2000 | |
| Applications | Tested Applications: WB, ELISA Cited Applications: | WB : Daudi | re Controls: audi cells, J774A.1 cells, RAW 264.7 cells, mouse ssue, rat lung tissue | |
| | WB Species Specificity: human, mouse, rat Cited Species: human | | | |
| Background Information | Janus kinase 1(JAK1), is a member of a class of protein-tyrosine kinases (PTK) characterized by the presence of a second phosphotransferase-related domain immediately N-terminal to the PTK domain. JAK1 is essential for signaling for certain type I and type II cytokines. JAK1 plays a critical role in initiating responses to multiple major cytokine receptor families. Expression of JAK1 in cancer cells enables individual cells to contract, potentially allowing them to escape their tumor and metastasize to other parts of the body. | | | |
| Storage | Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50 | | | |
| *** 20ul sizes contain 0.1% BSA | Aliquoting is unnecessary for -20 $^{\circ}$ C s | torage | | |

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data

Daudi JTAA RAW 264.7 mouse who 250 kDa→ 150 kDa-100 kDa-70 kDa→ 50 kDa→ 40 kDa→ 30 kDa→ 20 kDa→

Various lysates were subjected to SDS PAGE followed by western blot with 27284-1-AP (JAK1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.