

For Research Use Only

MMP-9 (Middle) Polyclonal antibody

Catalog Number: 27306-1-AP

65 Publications



Basic Information

Catalog Number:

27306-1-AP

Size:

150ul, Concentration: 350 ug/ml by Nanodrop and 267 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG26132

GenBank Accession Number:

BC006093

GeneID (NCBI):

4318

UNIPROT ID:

P14780

Full Name:

matrix metalloproteinase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase)

Calculated MW:

707 aa, 78 kDa

Observed MW:

92 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:500-1:1000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:50-1:500

Applications

Tested Applications:

WB, IHC, IP, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse

Cited Species:

human, mouse, rat

Positive Controls:

WB : human saliva tissue,

IP : human saliva tissue,

IHC : human tonsillitis tissue, human lung cancer tissue, human spleen tissue, mouse liver tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, tissue remodeling, and disease processes, such as arthritis or metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. Matrix metalloproteinase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase) (MMP9, synonyms: GELB, CLG4B) degrades collagens type IV and V. Studies in rhesus monkeys suggest that MMP9 is involved in IL-8-induced mobilization hematopoietic progenitor cells from bone marrow, and murine studies suggest a role in tumor-associated tissue remodeling. The pro-MMP9 is 92 kDa, and it can be detected a processed form of 68 kDa or 82 kDa. This protein can exist as a dimer of 180 kDa (PMID:7492685).

Notable Publications

Author	Pubmed ID	Journal	Application
XIAOYUE FENG	34528694	Oncol Rep	WB
WANG Xiao-He	34688464	Chin J Nat Med	WB
Cong Xu	34868365	Oncol Lett	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human saliva were subjected to SDS PAGE followed by western blot with 27306-1-AP (MMP9 (Middle) antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




IP result of anti-MMP9 (Middle) (IP:27306-1-AP, 4ug; Detection:27306-1-AP 1:300) with human saliva lysate 800 ug.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 27306-1-AP (MMP9 (Middle) antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 27306-1-AP (MMP9 (Middle) antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 27306-1-AP (MMP9 (Middle) antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).