## For Research Use Only

## B7H6 Polyclonal antibody

Catalog Number: 27428-1-AP



**Purification Method:** 

WB 1:1000-1:4000

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number:

GenBank Accession Number:

BC136800

GeneID (NCBI):

150ul, Concentration: 450 ug/ml by

374383

Nanodrop: Rabbit

27428-1-AP

**UNIPROT ID:** Q68D85

Isotype:

Full Name: hypothetical protein DKFZp686024166

IgG Immunogen Catalog Number:

Calculated MW:

AG26532

51 kDa

Observed MW: 75-80 kDa

**Applications** 

**Tested Applications:** 

WB, ELISA

Species Specificity:

human

Positive Controls:

WB: HEK-293 cells, HeLa cells, HepG2 cells

## **Background Information**

B7-H6, also known as NCR3LG1, is a member of the B7 family of immunoreceptors. It is a 454-aa-long type I transmembrane protein consisting of an extracellular region, a transmembrane segment, and a cytoplasmic domain which contains an immunoreceptor tyrosine-based inhibitory motif (ITIM), an SH2-binding domain, and an SH3binding motif. B7-H6 triggers NKp30-mediated activation of human NK cells (PMID: 19528259). It has been reported that B7H6 is expressed in various types of tumor cells and plays a prominent role in natural killer cell immunosurveillance (PMID: 35697295). The apparent molecular weight of B7-H6 is larger than the calculated molecular weight of 51 kDa due to glycosylation (PMID: 19528259).

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 27428-1-AP (B7H6 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.