

For Research Use Only

TRIM8 Polyclonal antibody

Catalog Number: 27463-1-AP

Featured Product

6 Publications



Basic Information

Catalog Number:

27463-1-AP

Size:

150ul, Concentration: 800 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG26517

GenBank Accession Number:

BC021925

GeneID (NCBI):

81603

UNIPROT ID:

Q9BZR9

Full Name:

tripartite motif-containing 8

Calculated MW:

61 kDa

Observed MW:

61 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF, IP

Species Specificity:

human

Cited Species:

human

Positive Controls:

WB : HepG2 cells, L02 cells

IHC : human breast cancer tissue, human liver cancer tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

TRIM8 (Tripartite motif containing protein 8), also known as RNF27 (RING finger protein 27), a member of TRIM family proteins, is known to play a dual role as both tumor suppressor and oncogene, and to function at the crosstalk of cancer and innate immunity. TRIM8 can function at two subcellular sites-nucleus and cytoplasm-to regulate NF- κ B, one of the central signaling pathways that plays a critical role in carcinogenesis and inflammatory diseases (PMID: 33230447, 22262183). TRIM8 is composed of 551 amino acids (aa) with a molecular weight of 61.5 kDa, contains a Nuclear Localization Signal (NLS), and forms specific nuclear structures similar to the TRIM19/PML Nuclear Bodies (NBs) (PMID: 33807506, 29182544).

Notable Publications

Author	Pubmed ID	Journal	Application
Lei Li	33129786	Exp Mol Pathol	WB
Dachang Liu	35102149	Sci Rep	WB
Mingxue Zhu	39039652	Adipocyte	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




L02 cells were subjected to SDS PAGE followed by western blot with 27463-1-AP (TRIM8 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



HepG2 cells were subjected to SDS PAGE followed by western blot with 27463-1-AP (TRIM8 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 27463-1-AP (TRIM8 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 27463-1-AP (TRIM8 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).