

For Research Use Only

ECSIT Polyclonal antibody, PBS Only

Catalog Number: 27548-1-PBS



Basic Information

Catalog Number:

27548-1-PBS

Size:

100ug, Concentration: 1 mg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG26877

GenBank Accession Number:

BC008279

GeneID (NCBI):

51295

UNIPROT ID:

Q9BQ95

Full Name:

ECSIT homolog (Drosophila)

Calculated MW:

49 kDa

Observed MW:

49 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, IHC, Indirect ELISA

Species Specificity:

human, mouse

Background Information

ECSIT is a cytosolic adaptor protein associated with the toll-like receptor pathway. It has a distinct N-terminal mitochondrial targeting sequence, pentatricopeptide repeat motif, and a C-terminal pleckstrin homology domain. ECSIT regulates many biological processes like embryonic development, inflammation, cardiac function, and assembly of mitochondrial complex I. In macrophages, the activation of TLRs by inflammatory signals promotes the translocation of TRAF6 to mitochondria where it engages ECSIT to stimulate the production of mitochondrial reactive oxygen species (ROS).

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human ovary cancer tissue slide using 27548-1-AP (ECSIT antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 27548-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human ovary cancer tissue slide using 27548-1-AP (ECSIT antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 27548-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 27548-1-AP (ECSIT antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 27548-1-PBS in a different storage buffer formulation.