For Research Use Only

DBNDD2 Polyclonal antibody

Catalog Number: 27623-1-AP 1 Publications



Basic Information

Catalog Number: GenBank Accession Number:

27623-1-AP BC001105
Size: GeneID (NCBI):

150ul , Concentration: 1000 ug/ml by 55861
Nanodrop and 467 ug/ml by Bradford method using BSA as the standard; Q9BQY9
Source: Full Name:

Rabbit dysbindin (dystrobrevin binding lsotype: protein 1) domain containing 2

IgG Calculated MW:
Immunogen Catalog Number: 261 aa, 28 kDa
AG26356 Observed MW:

28-29 kDa

Applications

Tested Applications: WB, IHC, ELISA

Cited Applications:

WB

Species Specificity: Human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: SH-SY5Y cells, mouse brain tissue, rat brain tissue

Purification Method:

WB 1:1000-1:4000

IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: rat brain tissue,

Notable Publications

 Author
 Pubmed ID
 Journal
 Application

 Lea Elsholz
 34831053
 Cells
 WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



mouse and rat brain tissues were subjected to SDS PAGE followed by western blot with 27623-1-AP (DBNDD2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 27623-1-AP (DBNDD2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 27623-1-AP (DBNDD2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 27623-1-AP (DBNDD2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).