

For Research Use Only

Nexilin Polyclonal antibody

Catalog Number: 27737-1-AP



Basic Information

Catalog Number: 27737-1-AP	GenBank Accession Number: BC114444	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 650 µg/ml by Nanodrop;	GeneID (NCBI): 91624	Recommended Dilutions: WB 1:20000-1:100000 IHC 1:50-1:500 IF 1:50-1:500
Source: Rabbit	Full Name: nexilin (F actin binding protein)	
Isotype: IgG	Calculated MW: 675 aa, 81 kDa	
Immunogen Catalog Number: AG26938	Observed MW: 80-85 kDa	

Applications

Tested Applications:

IF, IHC, WB, ELISA

Species Specificity:

Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse heart tissue, mouse skeletal muscle tissue, rat heart tissue

IHC : mouse heart tissue,

IF : mouse heart tissue,

Background Information

Nexilin, Also known as NEXN, NELIN, is a filamentous actin-binding protein that localizes to focal contacts and may be involved in cell adhesion and migration (PMID: 15823560). Nexilin is abundantly expressed in heart and skeletal muscle and has an essential role in the maintenance of Z line and sarcomere integrity (PMID: 12053183, 19881492). Mutations in Nexilin have been associated with cardiomyopathy dilated type 1CC (CMD1CC) and cardiomyopathy familial hypertrophic type 20 (CMH20) (PMID: 19881492, 20970104).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 27737-1-AP (Nexilin antibody) at dilution of 1:80000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 27737-1-AP (Nexilin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 27737-1-AP (Nexilin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse heart tissue using Nexilin antibody (27737-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse heart tissue using Nexilin antibody (27737-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).