

For Research Use Only

PHLDB2 Polyclonal antibody

Catalog Number: 27940-1-AP



Basic Information

Catalog Number:

27940-1-AP

Size:

150ul, Concentration: 800 ug/ml by Nanodrop and 433 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG27331

GenBank Accession Number:

BC142678

GeneID (NCBI):

90102

UNIPROT ID:

Q86SQ0

Full Name:

pleckstrin homology-like domain, family B, member 2

Calculated MW:

1253 aa, 142 kDa

Observed MW:

150 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:8000

IHC 1:400-1:1600

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : A431 cells,

IHC : human skin cancer tissue, human placenta tissue, mouse heart tissue

Background Information

The pleckstrin homology-like domain (PHLD) class of proteins are multifunctional proteins comprised of two families of proteins, PHLDA and PHLDB, each with 3 members: PHLDA1, PHLDA2, PHLDA3, PHLDB1, PHLDB2, and PHLDB3 (PMID: 32429563). All the family members possess a pleckstrin homology (PH) domain. PHLB2, also known as LL5-beta, contains two predicted coiled-coil domains, a serine-rich region predicted to contain multiple protein kinase C phosphorylation sites in addition to a PH domain (PMID: 15851520). PHLB2 is a filamin-binding protein that is an effector of the PI3K signaling pathway (PMID: 17174070). It is a PtdIns(3,4,5)P3 (PIP3) binding protein and can translocate to the plasma membrane in a receptor and PI3K-dependant manner (PMID: 12376540). PHLB2 has also been reported as a component of the postsynaptic apparatus and a regulator of postsynaptic differentiation (PMID: 15851520).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




A431 cells were subjected to SDS PAGE followed by western blot with 27940-1-AP (PHLDB2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded skin cancer slide using 27940-1-AP (PHLDB2 antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded skin cancer slide using 27940-1-AP (PHLDB2 antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 27940-1-AP (PHLDB2 antibody) at dilution of 1:600 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 27940-1-AP (PHLDB2 antibody) at dilution of 1:600 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).