For Research Use Only

CD19 Polyclonal antibody

Catalog Number: 27949-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number: 27949-1-AP

GenBank Accession Number:

Purification Method: Antigen affinity purification

BC006338 GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 267 µg/ml by

930

95 kDa

WB 1:1000-1:8000 IHC 1:50-1:1000

IF 1:50-1:500

Bradford method using BSA as the standard:

Full Name: CD19 molecule

Calculated MW:

556 aa, 61 kDa

Observed MW:

Immunogen Catalog Number:

AG27656

Rabbit

Isotype:

IgG

Applications

Tested Applications:

IF, IHC, WB, ELISA

Cited Applications:

IHC. WB

IHC: human tonsillitis tissue, mouse colon tissue,

Species Specificity:

human lymphoma tissue, rat colon tissue, rat thymus tissue

Human, Rat, Mouse

IF: Raji cells,

Positive Controls:

WB: Daudi cells, Raji cells

Cited Species: human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

CD19 is a B cell-specific molecule that controls B cell activation by complexing with the BCR. CD19 is a member of the Ig superfamily and is the dominant component for the signaling complex on B cells that includes CD21, CD81, and CD22. Humans with CD19 deficiency have reduced proliferative responses to BCR stimulation in vitro and mount impaired Ab responses to vaccination.

Notable Publications

Author	Pubmed ID	Journal	Application
Hirofumi Watanabe	36609460	Inflamm Regen	IHC
Changwei Zhang	38244740	Int J Biol Macromol	WB
Chen Su	37629071	Int J Mol Sci	IHC

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




WB result of CD19 antibody (27949-1-AP; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CD19 transfected Raji cells.




Daudi cells were subjected to SDS PAGE followed by western blot with 27949-1-AP (CD19 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 27949-1-AP (CD19 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 27949-1-AP (CD19 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Raji cells using CD19 antibody (27949-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).