

For Research Use Only

Ki67 Polyclonal antibody

Catalog Number: 28074-1-AP **29 Publications**



Basic Information

Catalog Number: 28074-1-AP	GenBank Accession Number: NM_001081117	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 400 µg/ml by Nanodrop;	GeneID (NCBI): 17345	Recommended Dilutions: IHC 1:1000-1:4000 IF 1:400-1:1600
Source: Rabbit	UNIPROT ID: E9PVX6	
Isotype: IgG	Full Name: antigen identified by monoclonal antibody Ki 67	
Immunogen Catalog Number: AG27894	Calculated MW: 351 kDa	

Applications

Tested Applications: IF, FC, IHC, ELISA	Positive Controls: IHC : mouse spleen tissue, IF : NIH/3T3 cells, mouse colon tissue, mouse spleen tissue
Cited Applications: WB, IF, IHC, Cell treatment	
Species Specificity: mouse	
Cited Species: human, rat, mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

The Ki-67 protein (also known as MKI67) is a cellular marker for proliferation. Ki67 is present during all active phases of the cell cycle (G1, S, G2 and M), but is absent in resting cells (G0). Cellular content of Ki-67 protein markedly increases during cell progression through S phase of the cell cycle. Therefore, the nuclear expression of Ki67 can be evaluated to assess tumor proliferation by immunohistochemistry.

Notable Publications

Author	Pubmed ID	Journal	Application
Wenbing Wu	36147440	J Oncol	IHC
Keiko Haraguchi-Suzuki	36074048	Genes Cells	WB,IF
Zifan Wang	36203070	In Vitro Cell Dev Biol Anim	IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 28074-1-AP (ki67 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 28074-1-AP (ki67 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using ki67 antibody (28074-1-AP) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of unfixed frozen OCT-embedded mouse spleen tissue using ki67 antibody (28074-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Immunofluorescent analysis of unfixed frozen OCT-embedded mouse spleen tissue using ki67 antibody (28074-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1×10^6 NIH/3T3 cells were intracellularly stained with 0.2 μ g Anti-Mouse ki67 (28074-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 μ g Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).