For Research Use Only

Caspase 14 Polyclonal antibody Catalog Number: 28136-1-AP




Basic Information	Catalog Number: 28136-1-AP	GenBank Accession Number: BC069541	Purification Method: Antigen affinity purification		
	Size: 150ul, Concentration: 800 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG27242	GenelD (NCBI): 23581	Recommended Dilutions: WB 1:1000-1:6000		
		UNIPROT ID:			
		P31944 Full Name: caspase 14, apoptosis-related cysteine peptidase Calculated MW: 242 aa, 28 kDa			
				Observed MW: 35 kDa	
				Applications	Tested Applications:
		WB, ELISA Species Specificity:			
	human, mouse, rat				
Background Information	Caspase14 belongs to the caspase fa role in apoptosis. Caspase14 exist as small subunits (p20 and p10) at Ile15 Caspase14 plays a role in protein ma has been detected in several of epith carcinogenesis and disease progress indicating that caspase14 may be an	inactive proenzymes that undergo 52/Lys153 residues, which is different aturation of filaggrin, and possibly i lelial malignancies, indicating that ion of cancers. Caspase14 can binds anti-apoptotic protein. In the LADC esent Caspase14 and the phosphon	n DNA repair. Overexpression of Caspase14 Caspase14 could play a role in 5 the apoptosis-inducing factor (AIF) specimens, two bands can be detected with ylated Caspase14(pCasp-14) respectively.		
Background Information	Caspase 14 belongs to the caspase fa role in apoptosis. Caspase 14 exist as small subunits (p20 and p10) at Ile 15 Caspase 14 plays a role in protein ma has been detected in several of epith carcinogenesis and disease progress indicating that caspase 14 may be an MW of 35 kDa and 37 kDa which repr (PMID:21567094, 12200134, 1020365	inactive proenzymes that undergo 52/Lys153 residues, which is different turation of filaggrin, and possibly in elial malignancies, indicating that ion of cancers. Caspase14 can binds anti-apoptotic protein. In the LADC esent Caspase14 and the phosphory 28, 17515931, 16061209). This antic eter shipment.	proteolytic processing to produce large and int to other caspases (Asp residue). In DNA repair. Overexpression of Caspase14 Caspase14 could play a role in is the apoptosis-inducing factor (AIF) specimens, two bands can be detected with ylated Caspase14(pCasp-14) respectively.		

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28136-1-AP (Caspase 14 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.