

For Research Use Only

CRLF2 Polyclonal antibody

Catalog Number: 28365-1-AP



Basic Information

Catalog Number: 28365-1-AP	GenBank Accession Number: NM_001012288	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 350 ug/ml by Nanodrop;	GeneID (NCBI): 64109	Recommended Dilutions: WB: 1:500-1:2000
Source: Rabbit	UNIPROT ID: Q9HC73	
Isotype: IgG	Full Name: cytokine receptor-like factor 2	
Immunogen Catalog Number: AG26842	Calculated MW: 42 kDa	
	Observed MW: 42 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HepG2 cells, Jurkat cells, LO2 cells
Species Specificity: human	

Background Information

Cytokine receptor-like factor 2 (CRLF2), also known as the thymic stromal derived lymphopoietin receptor (TSLPR), which in combination with the interleukin 7 receptor α chain (IL-7R) forms the receptor for thymic stromal lymphopoietin. CRLF2 is a type I cytokine receptor. CRLF2 rearrangements occur either as a translocation to the immunoglobulin heavy-chain enhancer region (IGH-CRLF2) or by a deletion of upstream PAR1 that leads to joining of CRLF2 to adjacent P2RY8. CRLF2 is expressed in heart, skeletal muscle, kidney and adult and fetal liver and is primarily expressed in dendrites and monocytes. (PMID: 20807819; 31644323; 33485429)

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28365-1-AP (CRLF2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



WB result of CRLF2 antibody (28365-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-CRLF2 transfected HepG2 cells.