For Research Use Only

BOP1 Polyclonal antibody

Catalog Number: 28366-1-AP

2 Publications



Basic Information

Catalog Number: 28366-1-AP

GenBank Accession Number:

Purification Method: Antigen affinity purification

Size:

BC013787 GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 300 µg/ml by

23246

WB 1:500-1:2000

Nanodrop and 200 µg/ml by Bradford Full Name:

block of proliferation 1

IHC 1:50-1:500 IF 1:50-1:500

method using BSA as the standard;

Calculated MW:

Rabbit Isotype:

84 kDa Observed MW:

IgG

117 kDa

Immunogen Catalog Number:

AG26669

Applications

Tested Applications:

IF, IHC, WB, ELISA

Cited Applications:

WB: DU 145 cells, NCI-H1299 cells, HeLa cells

IHC: human colon cancer tissue, mouse testis tissue

IF: PC-3 cells, MCF-7 cells

Positive Controls:

Species Specificity: Human, Mouse **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Block of Proliferation 1(BOP1) protein was first isolated from a mouse, and the truncation protein may led to cell cycle arrest. Bop1 plays a role in processing of the 28S and 5.8S rRNAs (ribosomal RNAs) to regulate ribosome biogenesis. And Bop1 also act as a component of Pes1/Bop1/WDR21 (PeBoW) complex for mature 605 ribosome, and consequently acts during cell division at G1 checkpoints (PMID:18670790; PMID:26940494). What's more, abnormal expression of BOP1 was associated with liver or colorectal cancers(PMID:16804918). The molecular weight of the protein we detected was slightly higher, about 117 kd, which is also seen in the article (PMID: 30100995).

Notable Publications

Author	Pubmed ID	Journal	Application
Lanjuan Mi	34868955	Front Oncol	WB
Miao Bai	34512161	Int J Biol Sci	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 28366-1-AP (BOP1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed PC-3 cells using 28366-1-AP (BOP1 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 28366-1-AP (BOP1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 28366-1-AP (BOP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).