For Research Use Only

## ATG7 Polyclonal antibody Catalog Number: 28413-1-AP Featured Product




Basic Information	Catalog Number: 28413-1-AP	GenBank Accession Number: BC000091	Purification Method: Antigen affinity purification
	Size: 150ul , Concentration: 500 ug/ml by Nanodrop; Source: Rabbit	GenelD (NCBI): 10533	Recommended Dilutions: WB 1:1000-1:5000 IHC 1:50-1:500
		UNIPROT ID: O95352	
		Full Name:	
	lsotype: IgG	ATG7 autophagy related 7 homolog (S. cerevisiae)	
	Immunogen Catalog Number: AG29172	Calculated MW: 78 kDa	
		Observed MW: 68-70 kDa	
Applications	Tested Applications: WB, IHC, ELISA Species Specificity: human	Positive Controls:	
		WB : HeLa cells, IHC : human cervical cancer tissue,	
	Note-IHC: suggested antigen i TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen	
Background Information	Atg7 is an E1-like enzyme that is specifically involved in autophagosome formation and is essential for autophagy. As an autophagic-related protein it is required for linking to Atg12, Atg5 and Atg8, which are essential for Atg conjugation and autophagosome formation. Atg7 has been reported as an important regulator of autophagy with starvation-induced or chemotherapeutic agent treatment. The high expression level of ATG7 is related to the survival of patients with breast cancer. There are several isoforms of ATG7 protein ranged from 68 kDa to 78 kDa.		
Storage	Storage: Store at -20°C. Stable for one year aff Storage Buffer: PBS with 0.02% sodium azide and 50		
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for $-20^{\circ}$ C s	•••	


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



HeLa cell lysates were subjected to SDS PAGE followed by western blot with 28413-1-AP (ATG7 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



WB result of ATG7 antibody (28413-1-AP; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ATG7 transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 28413-1-AP (ATG7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).