

For Research Use Only

NCX1 Polyclonal antibody

Catalog Number: 28447-1-AP **1 Publications**



Basic Information

Catalog Number: 28447-1-AP	GenBank Accession Number: NM_021097	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 600 µg/ml by Nanodrop;	GeneID (NCBI): 6546	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:200-1:1600
Source: Rabbit	Full Name: solute carrier family 8 (sodium/calcium exchanger), member 1	
Isotype: IgG	Calculated MW: 109 kDa	
Immunogen Catalog Number: AG28952	Observed MW: 70 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls:
Cited Applications: WB	WB : mouse heart tissue, rat heart tissue, rat kidney tissue
Species Specificity: Human, mouse, rat	IHC : human kidney tissue,
Cited Species: mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Hao Wu	36473990	Acta Pharmacol Sin	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 28447-1-AP (NCX1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 28447-1-AP (NCX1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 28447-1-AP (NCX1 antibody) at dilution of 1:1600 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).