

For Research Use Only

# NMDAR2A/GRIN2A Polyclonal antibody



Catalog Number: 28525-1-AP

6 Publications

## Basic Information

<b>Catalog Number:</b> 28525-1-AP	<b>GenBank Accession Number:</b> NM_000833	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul, Concentration: 780 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 2903	<b>Recommended Dilutions:</b> WB 1:1000-1:6000 IP 0.5-4.0 ug for IP and 1:1000-1:8000 for WB
<b>Source:</b> Rabbit	<b>Full Name:</b> glutamate receptor, ionotropic, N-methyl D-aspartate 2A	<b>IHC 1:50-1:500</b>
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 165 kDa	
<b>Immunogen Catalog Number:</b> AG29101	<b>Observed MW:</b> 160-180 kDa	

## Applications

**Tested Applications:**  
IHC, IP, WB, ELISA

**Cited Applications:**  
IF, WB

**Species Specificity:**  
mouse, rat

**Cited Species:**  
human, rat, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

**Positive Controls:**

**WB:** mouse brain tissue, rat brain tissue

**IP:** rat brain tissue,

**IHC:** mouse brain tissue,

## Background Information

GRIN2A (glutamate ionotropic receptor NMDA type subunit 2A), also known as NMDAR2A. And its molecular weight is 165 kDa. GRIN2A is located in cell projection, dendritic spine, cell membrane, synapse, postsynaptic cell membrane, cytoplasmic vesicle membrane, which is expressed in many tissues, highest expression in brain and heart. This gene encodes a member of the glutamate-gated ion channel protein family. The encoded protein is an N-methyl-D-aspartate (NMDA) receptor subunit. NMDA receptors are both ligand-gated and voltage-dependent, and are involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. These receptors are permeable to calcium ions, and activation results in a calcium influx into post-synaptic cells, which results in the activation of several signaling cascades. Disruption of this gene is associated with focal epilepsy and speech disorder with or without cognitive disability. Alternative splicing results in multiple transcript variants.

## Notable Publications

Author	Pubmed ID	Journal	Application
Kangyu Jin	36103758	Psychiatry Res	WB
Jie Du	36483743	Front Pharmacol	WB
XiaoHuan Liu	35340131	Andrology	WB

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 28525-1-AP (NMDAR2A/GRIN2A antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 28525-1-AP (NMDAR2A/GRIN2A antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 28525-1-AP (NMDAR2A/GRIN2A antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-NMDAR2A/GRIN2A(IP:28525-1-AP, 4ug; Detection:28525-1-AP 1:4000) with rat brain tissue lysate 1120 ug.