

For Research Use Only

# SUPT16H Polyclonal antibody

Catalog Number: 28598-1-AP

Featured Product



## Basic Information

<b>Catalog Number:</b> 28598-1-AP	<b>GenBank Accession Number:</b> NM_007192	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 600 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 11198	<b>Recommended Dilutions:</b> WB 1:2000-1:16000 IF 1:200-1:800
<b>Source:</b> Rabbit	<b>Full Name:</b> suppressor of Ty 16 homolog (S. cerevisiae)	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 120 kDa	
<b>Immunogen Catalog Number:</b> AG29459	<b>Observed MW:</b> 120 kDa	

## Applications

**Tested Applications:**  
IF, WB, ELISA

**Species Specificity:**  
Human, mouse, rat

### Positive Controls:

**WB:** A549 cells, MCF-7 cells, HeLa cells, mouse brain tissue, rat brain tissue

**IF:** HeLa cells,

## Background Information

SUPT16H, also named as FACT140, FACTP140, SPT16 and CDC68, belongs to the peptidase M24 family and SPT16 subfamily. SUPT16H is a component of the FACT complex, a general chromatin factor that acts to reorganize nucleosomes. The FACT complex is involved in multiple processes that require DNA as a template such as mRNA elongation, DNA replication and DNA repair. During transcription elongation the FACT complex acts as a histone chaperone that both destabilizes and restores nucleosomal structure. It facilitates the passage of RNA polymerase II and transcription by promoting the dissociation of one histone H2A-H2B dimer from the nucleosome, then subsequently promotes the reestablishment of the nucleosome following the passage of RNA polymerase II. The FACT complex is probably also involved in phosphorylation of 'Ser-392' of p53/TP53 via its association with CK2 (casein kinase II). It also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene. The antibody is specific to SUPT16H.

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28598-1-AP (SUPT16H antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



WB result of SUPT16H antibody (28598-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SUPT16H transfected MCF-7 cells.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using SUPT16H antibody (28598-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).