For Research Use Only

SUPT16H Polyclonal antibody, PBS Only 🔽

Catalog Number:28598-1-PBS

Featured Product

Antibodies | ELISA kits | Proteins www.ptglab.com

## **Basic Information**

Catalog Number: 28598-1-PBS Size: 100ug , Concentration: 1 mg/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG29459

GenBank Accession Number: NM\_007192 GeneID (NCBI): 11198 UNIPROT ID: Q9Y5B9 Full Name: suppressor of Ty 16 homolog (S. cerevisiae) Calculated MW: 120 kDa Observed MW: 120 kDa

#### Purification Method: Antigen affinity purification

Applications

Tested Applications: WB, IHC, IF/ICC, Indirect ELISA Species Specificity: human, mouse, rat

#### **Background Information**

SUPT16H, also named as FACT140, FACTP140, SPT16 and CDC68, belongs to the peptidase M24 family and SPT16 subfamily. SUPT16H is a component of the FACT complex, a general chromatin factor that acts to reorganize nucleosomes. The FACT complex is involved in multiple processes that require DNA as a template such as mRNA elongation, DNA replication and DNA repair. During transcription elongation the FACT complex acts as a histone chaperone that both destabilizes and restores nucleosomal structure. It facilitates the passage of RNA polymerase II and transcription by promoting the dissociation of one histone H2A-H2B dimer from the nucleosome, then subsequently promotes the reestablishment of the nucleosome following the passage of RNA polymerase II. The FACT complex is probably also involved in phosphorylation of 'Ser-392' of p53/TP53 via its association with CK2 (casein kinase II). It also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transcrepression of the CYP27B1 gene. The antibody is specific to SUPT16H.


### Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

# Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 28598-1-AP (SUPT16H antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 28598-1-PBS in a different storage buffer formulation.



WB result of SUPT16H antibody (28598-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SUPT16H transfected MCF-7 cells. This data was developed using the same antibody clone with 28598-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 28598-1-AP (SUPT16H antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 28598-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using SUPT16H antibody (28598-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 28598-1-PBS in a different storage buffer formulation.