For Research Use Only

Synaptotagmin-3 Polyclonal antibody

Catalog Number: 28667-1-AP



Basic Information

Catalog Number:

28667-1-AP BC028379

Size: GenelD (NCBI):

150ul, Concentration: 800 ug/ml by 84258

Nanodrop; UNIPROT ID:

Source: Q9BQG1 Rabbit Full Name:

 Isotype:
 synaptotagmin III

 IgG
 Calculated MW:

 Immunogen Catalog Number:
 590 aa, 63 kDa

AG30027 Observed MW:

70-75 kDa

GenBank Accession Number:

Applications

Tested Applications:

WB, IHC, IF-P, ELISA

Species Specificity:

Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

IF-P 1:50-1:500

Purification Method:

WB 1:2000-1:12000 IHC 1:500-1:2000

Antigen affinity purification

Recommended Dilutions:

Positive Controls:

WB: mouse brain tissue, rat brain tissue

IHC: mouse brain tissue,
IF-P: mouse brain tissue.

Background Information

The synaptotagmins are integral membrane proteins of synaptic vesicles thought to serve as Ca(2+) sensors in the process of vesicular trafficking and exocytosis (PMID: 8058779). Synaptotagmin-3 (SYT3) is highly expressed in brain, adipose tissue, and heart (PMID: 30545844). Synaptotagmin-3 is highly enriched in synaptic plasma membranes (PMID: 10373432). It has been reported that synaptotagmin-3 internalizes AMPA receptors to depress synaptic strength and promote forgetting (PMID: 30545844).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




mouse brain tissue and rat brain tissue were subjected to SDS PAGE followed by western blot with 28667-1-AP (Synaptotagmin-3 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 28667-1-AP (Synaptotagmin-3 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 28667-1-AP (Synaptotagmin-3 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Synaptotagmin-3 antibody (28667-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).