For Research Use Only

SGLT2 Polyclonal antibody

Catalog Number: 28683-1-AP 2 Publications




Basic Information	Catalog Number: 28683-1-AP	GenBank Accession Number: BC131542	Purification Method: Antigen affinity purification	
	Size: 150ul, Concentration: 900 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG30120	GenelD (NCBI):	Recommended Dilutions:	
		5524 WB 1:500-1:2000		
		UNIPROT ID:	IHC 1:50-1:500 IF-P 1:50-1:500	
		1024		
		Full Name: solute carrier family 5		
		(sodium/glucose cotransporter),		
		member 2		
		Calculated MW: 672 aa, 73 kDa		
		Observed MW:		
		73 kDa		
Applications	Tested Applications:	HC, IF-P, ELISA WB : mouse kidney tissue, rat kidney tissue		
	WB, IHC, IF-P, ELISA			
	Cited Applications: WB			
	Species Specificity: Human, mouse, rat	IF-P : mouse kidney tissue, rat kidney tissue		
	Cited Species: mouse, rat			
	Note-IHC: suggested antigen ı TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen		
Background Information	transport glucose against the cell's ir (BBM) of epithelial cells in the early	nternal concentration gradient. Main segment of the proximal tubule, SGI nhibition of SGLT2 could improve gl	e electrochemical sodium gradient to ly expressed on brush border membrar LT2 mediates most of the glucose ucose homeostasis of diabetic patients	
	transport glucose against the cell's ir (BBM) of epithelial cells in the early reabsorption by the kidney overall. I which has been considered as a nove	nternal concentration gradient. Main segment of the proximal tubule, SGI nhibition of SGLT2 could improve gl	ly expressed on brush border membrar LT2 mediates most of the glucose	
Background Information	transport glucose against the cell's ir (BBM) of epithelial cells in the early reabsorption by the kidney overall. I which has been considered as a nove	nternal concentration gradient. Main segment of the proximal tubule, SGI nhibition of SGLT2 could improve gl al strategy for diabetes treatment.	ly expressed on brush border membrar LT2 mediates most of the glucose ucose homeostasis of diabetic patients	
	transport glucose against the cell's ir (BBM) of epithelial cells in the early reabsorption by the kidney overall. I which has been considered as a nove Author Put Tomoko Kodama 398	nternal concentration gradient. Main segment of the proximal tubule, SGI nhibition of SGLT2 could improve gl el strategy for diabetes treatment.	ly expressed on brush border membrar LT2 mediates most of the glucose ucose homeostasis of diabetic patients Application	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 28683-1-AP (SGLT2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 28683-1-AP (SGLT2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse kidney tissue using SGLT2 antibody (28683-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed rat kidney tissue using SGLT2 antibody (28683-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).