

For Research Use Only

SP7 Polyclonal antibody

Catalog Number: 28694-1-AP

2 Publications



Basic Information

Catalog Number:

28694-1-AP

Size:

150ul, Concentration: 350 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG29889

GenBank Accession Number:

NM_152860

GeneID (NCBI):

121340

UNIPROT ID:

Q8TDD2

Full Name:

Sp7 transcription factor

Calculated MW:

45 kDa

Observed MW:

45 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human

Cited Species:

human, rat

Positive Controls:

WB : Saos-2 cells,

Background Information

Sp7 is identified in several tissues, including bone, tooth, brain, and the reproductive tract (PMID: 11792318, 33135214, 36881265). In bone, Sp7 is expressed at multiple stages throughout the mesenchymal bone lineage and plays different roles in different osteocytes (PMID: 36881265). SP7 regulates osteoblast differentiation and bone formation through osteoblast target genes by recognizing AT-rich region. In osteocytes, SP7 regulates osteocyte maturation and intracortical remodeling through osteocyte target genes by recognizing TGA(G/T) TCA motif. SP7 also regulates chondrocyte differentiation and endochondral ossification (36881265).

Notable Publications

Author	Pubmed ID	Journal	Application
Taojing Yu	39956235	Int J Biol Macromol	WB
Lijun Li	39920313	Sci Rep	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28694-1-AP (SP7 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.