

For Research Use Only

# IFN gamma Polyclonal antibody

Catalog Number: 28722-1-AP



## Basic Information

<b>Catalog Number:</b> 28722-1-AP	<b>GenBank Accession Number:</b> BC070256	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 450 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 3458	<b>Recommended Dilutions:</b> WB 1:500-1:2000
<b>Source:</b> Rabbit	<b>Full Name:</b> IFN gamma	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 166 aa, 19 kDa	
<b>Immunogen Catalog Number:</b> AG30371	<b>Observed MW:</b> 20-25 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : PMA, Ionomycin and Brefeldin A treated NK-92 cells,
<b>Species Specificity:</b> Human	

## Background Information

Interferon gamma (IFNG) is a soluble cytokine that is the only member of the type II class of interferons. It is secreted by Th1 cells, cytotoxic T cells and NK cells. The cytokine is associated with antiviral, immunoregulatory and anti-tumor properties and is a potent activator of macrophages. It plays crucial roles in pathogen clearance. Aberrant IFNG expression is associated with a number of autoinflammatory and autoimmune diseases. It has been identified in many studies as a biomarker for pleural tuberculosis (TB). Mutations in this gene are associated with aplastic anemia.

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Non-treated, PMA, Ionomycin and Brefeldin A treated NK-92 cells were subjected to SDS PAGE followed by western blot with 28722-1-AP (IFNG antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.