

For Research Use Only

MDMX Polyclonal antibody

Catalog Number: 28747-1-AP



Basic Information

| | | |
|---|--|--|
| Catalog Number: 28747-1-AP | GenBank Accession Number: BC067299 | Purification Method: Antigen affinity purification |
| Size: 150ul , Concentration: 1000 µg/ml by Nanodrop; | GeneID (NCBI): 4194 | Recommended Dilutions: WB 1:2000-1:12000 IHC 1:50-1:500 |
| Source: Rabbit | Full Name: Mdm4 p53 binding protein homolog (mouse) | |
| Isotype: IgG | Calculated MW: 490 aa, 55 kDa | |
| Immunogen Catalog Number: AG30456 | Observed MW: 70-76 kDa | |

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A2780 cells, MCF-7 cells, MDA-MB-231 cells, U2OS cells

IHC: human ovary tumor tissue, human breast cancer tissue

Background Information

MDM4 is also named as MDMX. It inhibits the activity of the tumor suppressor p53 by primarily cooperating with the p53 feedback regulator MDM2. MDM4 resides mostly in the cytoplasm, but can be recruited to the nucleus by MDM2. (PMID:16511572). The predicted mass of the protein is 54 kDa, while the observed mass is 70-80 kDa, a difference which stated is probably due to phosphorylation or other posttranslational modification (PMID:9226370, 28097652). It can be ubiquitinated and degraded by MDM2 (PMID:16163388). It has 5 isoforms produced by alternative splicing.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 28747-1-AP (MDMX antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 28747-1-AP (MDM4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 28747-1-AP (MDM4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 28747-1-AP (MDMX antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).