For Research Use Only

MEF2A Polyclonal antibody

Catalog Number: 28819-1-AP



Basic Information

Catalog Number: GenBank Accession Number: 28819-1-AP BC013437

8819-1-AP BC013437 ize: GeneI D (NCBI):

150ul , Concentration: 350 µg/ml by 4205

Nanodrop; Full Name

Source: myocyte enhancer factor 2A

Rabbit Calculated MW:
Isotype: 499 aa, 54 kDa
IgG Observed MW:
Immunogen Catalog Number: 50-70kDa

AG30349

WB 1:500-1:2000 IHC 1:50-1:500

Purification Method:

Antigen affinity purification

Recommended Dilutions:

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human, mouse, rat

Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HUVEC cells, SH-SY5Y cells, THP-1 cells

IHC: rat brain tissue,

Background Information

MEF2A belongs to a family of DNA binding regulatory proteins. Each of these proteins which belong to the myocyte-specific enhancer factor-2 (MEF2) family binds to the MEF2 target DNA sequence present in the regulatory regions of many, if not all, muscle-specific genes. MEF2A plays diverse roles in the control of cell growth, survival and apoptosis via p38 MAPK signaling in muscle-specific and/or growth factor-related transcription. MEF2A also has key roles in cardiac and skeletal muscle development. MEF2A is especially abundant in granule neurons of the cerebellar cortex throughout the period of synaptogenesis. This antibody can detect two bands around 60 kDa to 70 kDa, the larger band may correspond to its post-translational modifications.(PMID: 15483225,16484498, 23486945)

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28819-1-AP (MEF2A antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 28819-1-AP (MEF2A antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).