For Research Use Only

## Phospho-SMAD1 (Ser187) Polyclonal antibody

www.ptglab.com

Catalog Number: 28865-1-AP

2 Publications

**Basic Information** 

Catalog Number:

28865-1-AP

Size:

100ul , Concentration: 350 ug/ml by

Nanodrop: Source:

Rabbit Isotype:

IgG

GenBank Accession Number: BC001878

GeneID (NCBI):

**UNIPROT ID:** Q15797

Full Name:

SMAD family member 1

Calculated MW:

52 kDa Observed MW: 60 kDa

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:4000

**Applications** 

**Tested Applications:** 

WB, ELISA

**Cited Applications:** 

WB

Species Specificity:

Human

**Cited Species:** 

rat

**Positive Controls:** 

WB: BMP2 treated HepG2 cells,

**Background Information** 

Transforming growth factor- $\beta$  (TGF- $\beta$ ) superfamily is recognized as one of the largest families of secreted multifunctional peptides exerting different biological effects on a large variety of cell types, such as regulation of hormone secretion, stimulation of extracellular matrix formation, the inhibition of proliferation of many cell types, cell survival, bone formation, and chemotaxis for inflammatory cells. One of the most important proteins that  $modulate\ TGF-\beta\ ligand\ activity\ is\ the\ SMAD\ family\ proteins.\ SMAD\ is\ one\ of\ the\ receptor-activated\ Smads.\ It's\ also$ a signal transducers of BMP signaling and binds to several proteins involved in ubiquitin-proteasome system (UPS).

**Notable Publications** 

| Author     | Pubmed ID | Journal                          | Application |
|------------|-----------|----------------------------------|-------------|
| You Peng   | 36398315  | Cardiol Res Pract                | WB          |
| Wanjing Li | 39411943  | Comb Chem High Throughput Screen | WB          |

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Non-treated HepG2 and BMP2 (HZ-1128) treated HepG2 cells were subjected to SDS PAGE followed by western blot with 28865-1-AP (Phospho-SMAD1 (Ser187) antibody) at dilution of 1:2000 incubated at 4°C overnight. The membrane was stripped and re-blotted with GAPDH antibody as loading control.