

For Research Use Only

Phospho-GRB10 (Ser476) Polyclonal antibody



Catalog Number: 28873-1-AP

Basic Information

Catalog Number: 28873-1-AP	GenBank Accession Number: BC024285	Purification Method: Antigen affinity purification
Size: 100ul , Concentration: 350 µg/ml by Nanodrop;	GeneID (NCBI): 2887	Recommended Dilutions: WB 1:500-1:2000
Source: Rabbit	Full Name: growth factor receptor-bound protein 10	
Isotype: IgG	Calculated MW: 536aa,61 kDa; 594aa,67 kDa	
	Observed MW: 70-80 kDa	

Applications

Tested Applications: WB,ELISA	Positive Controls: WB : λ phosphatase treated NIH/3T3 cells,
Species Specificity: Human, Mouse	

Background Information

Growth factor receptor-bound protein 10 (GRB10) is an adapter protein which modulates coupling of a number of cell surface receptor kinases with specific signaling pathways. GRB10 has three consensus domains including pleckstrin homology (PH) domain, SH2/SH3 domain and Ras-associating domain. By binding to kinases, GRB10 suppresses signals from activated receptors tyrosine kinases, including the INSR and INS-like growth factor (IGF1R) receptors. It may play a role in mediating INS-stimulated ubiquitination of INSR, leading to proteasomal degradation. It has been reported that GRB10 is a direct substrate of mechanistic/mammalian target of mTOR and can be phosphorylated at Ser150, Ser428 and Ser476 identified as MAPK-mediated phosphorylation sites. (PMID: 15952796, 24746805, 21659605)

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated NIH/3T3, phosphatase inhibitor treated and λ phosphatase treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 28873-1-AP (Phospho-GRB10 (Ser476) antibody) at dilution of 1:1000 incubated at 4°C overnight.