

For Research Use Only

Phospho-GRB10 (Ser476) Polyclonal antibody

Catalog Number: 28873-1-AP



Basic Information

Catalog Number:

28873-1-AP

Size:

100ul, Concentration: 350 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC024285

GeneID (NCBI):

2887

UNIPROT ID:

Q13322

Full Name:

growth factor receptor-bound protein 10

Calculated MW:

536aa, 61 kDa; 594aa, 67 kDa

Observed MW:

70-80 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human, Mouse

Positive Controls:

WB : λ phosphatase treated NIH/3T3 cells,

Background Information

Growth factor receptor-bound protein 10 (GRB10) is an adapter protein which modulates coupling of a number of cell surface receptor kinases with specific signaling pathways. GRB10 has three consensus domains including pleckstrin homology (PH) domain, SH2/SH3 domain and Ras-associating domain. By binding to kinases, GRB10 suppresses signals from activated receptors tyrosine kinases, including the INSR and INS-like growth factor (IGF1R) receptors. It may play a role in mediating INS-stimulated ubiquitination of INSR, leading to proteasomal degradation. It has been reported that GRB10 is a direct substrate of mechanistic/mammalian target of mTOR and can be phosphorylated at Ser150, Ser428 and Ser476 identified as MAPK-mediated phosphorylation sites. (PMID: 15952796, 24746805, 21659605)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated NIH/3T3, phosphatase inhibitor treated and λ phosphatase treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 28873-1-AP (Phospho-GRB10 (Ser476) antibody) at dilution of 1:1000 incubated at 4°C overnight.