

For Research Use Only

# Phospho-MEK6 (Ser207) Polyclonal antibody



Catalog Number: 28903-1-AP

## Basic Information

<b>Catalog Number:</b> 28903-1-AP	<b>GenBank Accession Number:</b> BC012009	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 100ul , Concentration: 600 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 5608	<b>Recommended Dilutions:</b> WB 1:1000-1:6000
<b>Source:</b> Rabbit	<b>Full Name:</b> mitogen-activated protein kinase kinase 6	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 334 aa, 37 kDa	
	<b>Observed MW:</b> 35-37 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : UV treated HEK-293 cells, HEK-293 cells, HeLa cells, Anisomycin treated HeLa cells
<b>Species Specificity:</b> Human	

## Background Information

MAPKK6 (Dual specificity mitogen activated protein kinase kinase 6), also known as MAP2K6 / MEK6 / MKK6, is a member of MAPKK protein kinase family. MKK6 plays an important role in intracellular signaling pathways leading toward activation of the p38 MAP kinase. MEK6 phosphorylates and activates p38 MAP kinase in response to inflammatory cytokines or environmental stress. Phosphorylation of Ser207 and Thr211 by MAP3Ks is critical for MEK6 activation, and acetylation of these residues by VopA blocks MEK6 activity (PMID: 26648936). The antibody also recognizes the ser218 phosphorylation site of MEK3.

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Non-treated HEK-293 cells, UV treated HEK-293 cells, non-treated HeLa cells and Anisomycin treated HeLa cells were subjected to SDS PAGE followed by western blot with 28903-1-AP (Phospho-MEK6 (Ser207) antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.