## Phospho-PRKD1 (Ser916) Polyclonal antibody

Catalog Number:28928-1-AP

2 Publications




Basic Information	Catalog Number: 28928-1-AP	GenBank Accession Number: NM_001330069	Purification Method: Antigen affinity purification	
	Size: 100ul , Concentration: 110 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GenelD (NCBI):	Recommended Dilutions:	
			WB 1:500-1:2000	
		UNIPROT ID: Q15139		
		Full Name:		
		protein kinase D1		
		Calculated MW: 102 kDa		
		Observed MW: 110 kDa		
Applications	Tested Applications: WB, ELISA	Positive Controls:		
	Cited Applications: WB	WB : Insu	lin treated NIH/3T3 cells,	
	Species Specificity: Human, Mouse			
	Cited Species: rat			
Background Information	cytoskeletal reorganization, Golgi fu heart, lung and other tissues. PKCs ha a PKD1 autophosphorylation site. PKI lipids, cross-linking of B- and T-cell m	KD1 is involved in cellular process nction, immune function and apop we been shown to regulate PKD1 D1 can be activated by growth fact eceptors and some G-protein coup	d of two cysteine-rich domains and a ses including protein secretion, proliferation otosis. It is widely expressed in thyroid, brai activation. It has been reported that ser 916 tors, oxidative stress, thrombin, bioactive oled receptors (GPCR). PKD1 is located mainl ombrane in activated cells. (PMID: 17306383	
	pleckstrin homology (PH) domain. PH cytoskeletal reorganization, Golgi fu heart, lung and other tissues. PKCs ha a PKD1 autophosphorylation site. PKI lipids, cross-linking of B- and T-cell r in the cytoplasm in unstimulated cel 24806360, 30101477, 21696630)	KD1 is involved in cellular process nction, immune function and apop we been shown to regulate PKD1 D1 can be activated by growth fact eceptors and some G-protein coup	es including protein secretion, proliferation ptosis. It is widely expressed in thyroid, brai activation. It has been reported that ser 916 tors, oxidative stress, thrombin, bioactive sled receptors (GPCR). PKD1 is located mainl	
	pleckstrin homology (PH) domain. PK cytoskeletal reorganization, Golgi fu heart, lung and other tissues. PKCs ha a PKD1 autophosphorylation site. PKI lipids, cross-linking of B- and T-cell rr in the cytoplasm in unstimulated cel 24806360, 30101477, 21696630) Author Put	KD1 is involved in cellular process nction, immune function and apop we been shown to regulate PKD1 D1 can be activated by growth fact eceptors and some G-protein coup Is, while PKD1 migrates to the me	es including protein secretion, proliferation boosis. It is widely expressed in thyroid, brai activation. It has been reported that ser 916 tors, oxidative stress, thrombin, bioactive oled receptors (GPCR). PKD1 is located mainl embrane in activated cells. (PMID: 17306383 Application	
Background Information	pleckstrin homology (PH) domain. Pk cytoskeletal reorganization, Golgi fu heart, lung and other tissues. PKCs ha a PKD1 autophosphorylation site. PKI lipids, cross-linking of B- and T-cell m in the cytoplasm in unstimulated cel 24806360, 30101477, 21696630) Author Put Yao Liu 333	KD1 is involved in cellular process nction, immune function and apop we been shown to regulate PKD1 of Cl can be activated by growth fact eceptors and some G-protein coup ls, while PKD1 migrates to the me omed ID Journal	es including protein secretion, proliferation ptosis. It is widely expressed in thyroid, brai activation. It has been reported that ser 916 tors, oxidative stress, thrombin, bioactive oled receptors (GPCR). PKD1 is located mainl mbrane in activated cells. (PMID: 17306383 Application icol WB	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Non-treated NIH/3T3 and Insulin treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 28928-1-AP (Phospho-PRKD1 (Ser916) antibody) at dilution of 1:1000 incubated at 4°C overnight. The membrane was stripped and re-blotted with GAPDH antibody as loading control.