

For Research Use Only

Phospho-MEK1 (Thr286) Polyclonal antibody

Catalog Number: 28933-1-AP



Basic Information

Catalog Number:

28933-1-AP

Size:

100ul, Concentration: 350 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC139729

GeneID (NCBI):

5604

ENSEMBL Gene ID:

ENSG00000169032

UNIPROT ID:

Q02750

Full Name:

mitogen-activated protein kinase kinase 1

Calculated MW:

43 kDa

Observed MW:

40-45 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human

Positive Controls:

WB : nocodazole treated HeLa cells,

Background Information

MAP2K1 encodes MAPK1, also known as MEK1. MEK1 variants can enhance MEK1 expression and ERK1 phosphorylation that together lead to continuous activation of MEK/ERK signaling pathway. MEK1 bind directly to ERK2 through a region in the N terminus of MEK. In addition, a proline-rich (PR) regulatory sequence in MEK is also involved in MEK-ERK association and signal propagation. The coupling between MEK1 and ERK2 is enhanced through phosphorylation on S298 in the MEK1 PR region, whereas phosphorylation on MEK1 T292 releases the complex. MEK1 enzymatic activity is regulated by site-specific phosphorylation that can be activated with phosphorylation of Ser217/Ser221 by Raf kinase or suppressed by phosphorylation of Thr286 and Thr292 by CDK1 and CDK5 or Thr292 and Thr386 by ERK1/2. (PMID: 31972311, PMID: 17928366, PMID: 22177953)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated HeLa cells and nocodazole treated HeLa cells were subjected to SDS PAGE followed by western blot with 28933-1-AP (Phospho-MEK1 (Thr286) antibody) at dilution of 1:3000 incubated at room temperature for 1 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.